


Florida Betony Control



Lawns-

1. February/March- Atrazine (safe on Centipede, St. Augustine, and dormant Bermuda)

Note the percentage-older Hi-Yield Atrazine contained 40%, newer contains 4%.

NEW-4%-Mix 4.3 oz in 1 gallon water into a pump sprayer, shake thoroughly-this will cover 500 square feet (50'x10')

Older-40%-Mix 8 oz of Atrazine with 8 oz. water into the Gilmour Sprayer, set dial on 3 (tablespoons), shake to mix the chemical with water and spray onto 5,450 square feet (54'x100')

2. April/May-Weed Out Broadleaf Control (granular not the spray-do not use on St. Augustine lawns –safe on Centipede, Bahia, Bermuda and Zoysia) Mow the lawn 1 to 2 days before application, water the lawn thoroughly at least 1 day before application to sustain watering until the next watering, Spread (10 lbs./5,000 sq. ft. when leaves are wet to hold the material and prevent dust drift, Do not wash from weed leaves for 1 to 2 days, Afterwards a thorough watering should be made. Reapply 20-30 days later if temperatures are not over 85 F.

3. October/November- Atrazine (safe on Centipede, St. Augustine, and dormant Bermuda) See instructions for February/March

Ornamentals-

1. Mix Killzall, Spreader Sticker and water in a clean plastic container (pail or bucket) 5 tablespoons of Killzall per gallon of water; 4 tablespoons of Hi-Yield Spreader Sticker per gallon of water, put on a water-proof glove (latex, plastic, rubber), then put on an absorbent cotton glove over it, wet the glove with Killzall solution, squeeze excess to prevent dripping, wet the leaves and stems of the weeds. Be careful in applying to wet only those plants you wish to kill! Killzall doesn't contaminate the soil.

2. Reapply on new shoots that emerge! Inspect every 6-8 weeks for re-emerging Florida Betony shoots!